

# Dog Owner Information about RIMADYL® Caplets / Chewable Tablets

CARPROFEN

Rimadyl® (pronounced “Rim-a-dill”) for Osteoarthritis and Post-Surgical Pain. Generic name: carprofen (“car-prō-fen”)

This summary contains important information about Rimadyl. You should read this information before you start giving your dog Rimadyl and review it each time the prescription is filled. This sheet is provided only as a summary and does not take the place of instructions from your veterinarian. Talk to your veterinarian if you do not understand any of this information or if you want to know more about Rimadyl.

## What is Rimadyl?

Rimadyl is a nonsteroidal anti-inflammatory drug (NSAID) that is used to reduce pain and inflammation (soreness) due to osteoarthritis and surgery in dogs. A licensed veterinarian must prescribe Rimadyl for your dog. It is available as a caplet and chewable tablet and is given to dogs by mouth.

Osteoarthritis (OA) is a painful condition caused by “wear and tear” of cartilage and other parts of the joints that may result in the following changes or signs in your dog:

- Limping or lameness
- Decreased activity or exercise (difficulty or reluctance in standing, climbing stairs, jumping or running)
- Stiffness or decreased movement of joints

To control surgical pain (for example, for surgeries such as spays, ear procedures or orthopedic repairs), your veterinarian may administer Rimadyl before the procedure and recommend that your dog be treated for several days after going home.

## What kind of results can I expect when my dog is on Rimadyl for OA?

While Rimadyl is not a cure for osteoarthritis, it can relieve the pain and inflammation of OA and improve your dog’s mobility.

- Response varies from dog to dog but can be quite dramatic.
- In most dogs, improvement can be seen in a matter of days.
- If Rimadyl is discontinued or not given as directed, your dog’s pain and inflammation may come back.

## Who should not take Rimadyl?

Your dog should not be given Rimadyl if he/she:

- Has had an allergic reaction to carprofen, the active ingredient of Rimadyl.
- Has had an allergic reaction to aspirin or other NSAIDs (for example, etodolac, meloxicam or deracoxib) such as hives, facial swelling, or red or itchy skin.

**Rimadyl should be given to dogs only.** Cats should not be given Rimadyl. Call your veterinarian immediately if your cat receives Rimadyl. People should not take Rimadyl. Keep Rimadyl and all medicines out of reach of children. Call your physician immediately if you accidentally take Rimadyl.

## How to give Rimadyl to your dog.

Rimadyl should be given according to your veterinarian’s instructions. Your veterinarian will tell you what amount of Rimadyl is right for your dog and for how long it should be given. Rimadyl Caplets should be given by mouth. Most dogs will take Rimadyl Chewable Tablets right out of your hand or the tablet can be placed in the mouth. Rimadyl may be given with or without food.

## What to tell/ask your veterinarian before giving Rimadyl.

*Talk to your veterinarian about:*

- The signs of OA you have observed (for example, limping, stiffness).
- The importance of weight control and exercise in the management of OA.
- What tests might be done before Rimadyl is prescribed.
- How often your dog may need to be examined by your veterinarian.
- The risks and benefits of using Rimadyl.

*Tell your veterinarian if your dog has ever had the following medical problems:*

- Experienced side effects from Rimadyl or other NSAIDs, such as aspirin
- Digestive upset (vomiting and/or diarrhea)
- Liver disease
- Kidney disease
- A bleeding disorder (for example, Von Willebrand’s disease)

*Tell your veterinarian about:*

- Any other medical problems or allergies that your dog has now or has had.
- All medicines that you are giving your dog or plan to give your dog, including those you can get without a prescription.

*Tell your veterinarian if your dog is:*

- Pregnant, nursing or if you plan to breed your dog.

## What are the possible side effects that may occur in my dog during Rimadyl therapy?

Rimadyl, like other drugs, may cause some side effects. Serious but rare side effects have been reported in dogs taking NSAIDs, including Rimadyl. Serious side effects can occur with or without warning and in rare situations result in death.

The most common NSAID-related side effects generally involve the stomach (such as bleeding ulcers). Kidneys and liver can also be affected. Look for the following side effects that can indicate your dog may be having a problem with Rimadyl or may have another medical problem:

- Decrease or increase in appetite
- Vomiting
- Change in bowel movements (such as diarrhea, or black, tarry or bloody stools)
- Change in behavior (such as decreased or increased activity level, incoordination, seizure or aggression)
- Yellowing of gums, skin, or whites of the eyes (jaundice)
- Change in drinking habits (frequency, amount consumed)
- Change in urination habits (frequency, color or smell)
- Change in skin (redness, scabs or scratching)

It is important to stop therapy and contact your veterinarian immediately if you think your dog has a medical problem or side effect from Rimadyl therapy. If you have additional questions about possible side effects, talk to your veterinarian.

## Can Rimadyl be given with other medicines?

Rimadyl should not be given with other NSAIDs (for example, aspirin, etodolac) or steroids (for example, cortisone, prednisone, dexamethasone, triamcinolone).

Tell your veterinarian about all medicines you have given your dog in the past, and any medicines that you are planning to give with Rimadyl. This should include other medicines that you can get without a prescription. Your veterinarian may want to check that all of your dog’s medicines can be given together.

## What do I do in case my dog eats more than the prescribed amount of Rimadyl?

Contact your veterinarian immediately if your dog eats more than the prescribed amount of Rimadyl.

## How to store Rimadyl Chewable Tablets.

Rimadyl Chewable Tablets are designed to taste good to animals. **Keep Rimadyl Chewable Tablets in a secured storage area out of the reach of your dog and other pets.** If your dog ingests more than your veterinarian prescribed, or if your other pets take Rimadyl Chewable Tablets, contact your veterinarian right away.

## What else should I know about Rimadyl?

This sheet provides a summary of information about Rimadyl. If you have any questions or concerns about Rimadyl or osteoarthritis or surgical pain, talk to your veterinarian.

As with all prescribed medicines, Rimadyl should only be given to the dog for which it was prescribed. It should be given to your dog only for the condition for which it was prescribed.

It is important to periodically discuss your dog’s response to Rimadyl at regular checkups. Your veterinarian will best determine if your dog is responding as expected and if your dog should continue receiving Rimadyl. To report a suspected adverse reaction call Zoetis at 1-888-963-8471.

Issued October 2014

Distributed by:

**zoetis**

Div. of Zoetis Inc  
Kalamazoo, MI 49007