

Betamethasone with Gentamicin Ophthalmic

(*bate-uh-meth-uh-sown* with *jen-ta-mye-sin*)

Category: Steroidal Anti-inflammatory Agent

Other Names for this Medication: Gentocin Durafil[®]

Common Dosage Forms: Veterinary: None. Formerly available as Gentocin Durafil[®]

as betamethasone 0.1% and gentamicin sulfate 0.3% ophthalmic suspension. **Human:** None. Although no longer commercially available, this combination product may be made (compounded) by qualified professionals.

This information sheet does not contain all available information for this medication. It is to help answer commonly asked questions and help you give the medication safely and effectively to your animal. If you have other questions or need more information about this medication, contact your veterinarian or pharmacist.

Key Information

- ▶ Shake well before using.
- ▶ Administer the drug properly to avoid contamination (ie, introducing bacteria into the bottle). Keep cap on bottle tightly closed when not in use.
- ▶ Wait 5 minutes after applying this medication before applying any other medications to the eye. If you are also giving an ophthalmic ointment, give the eye drops first to improve absorption of the medications.
- ▶ Store in the refrigerator or at controlled room temperature away from moisture and sunlight; do not freeze.
- ▶ Do not use in animals with an eye ulcer or with fungal or viral eye infections.
- ▶ Not for use in birds.

How is this medication useful?

Betamethasone with gentamicin ophthalmic solution is useful for treating inflammation of the eye caused by allergies, bacterial infections, or irritation from foreign objects in the eye. This medication is also useful in treating inflammation of the inside of the eye (uveitis). It may be used alone or with combinations of antibiotics when necessary. The FDA (U.S. Food & Drug Administration) has approved this drug for use in humans and in animals. You and your veterinarian can discuss why this drug is the most appropriate choice.

What should I tell my veterinarian to see if this medication can be safely given?

Many things might affect how well this drug will work in your animal. Be sure to discuss the following with your veterinarian so together you can make the best treatment decisions.

- ▶ Other drugs can interact with this medication, so be sure to tell your veterinarian and pharmacist what medications (including other eye medications, vitamins, supplements, or herbal therapies) you give your animal, including the amount and time you give each.
- ▶ Tell your veterinarian about any conditions or diseases your pet may have now or has had in the past.
- ▶ If your animal has been treated for the same disease or condition in the past, tell your veterinarian about the treatment and how well it did or didn't work.
- ▶ If your animal is pregnant or nursing, talk to your veterinarian about the risks of using this drug.

- ▶ Tell your veterinarian and pharmacist about any medication side effects (including allergic reactions, lack of appetite, diarrhea, itching, hair loss) your pet has developed in the past.

When should this medication not be used or be used very carefully?

No drug is 100% safe in all patients, but your veterinarian will discuss with you any specific concerns about using this drug in your animal.

This drug **SHOULD NOT** be used in:

- ▶ Animals that are allergic to betamethasone, gentamicin, or related drugs.
- ▶ Birds, as they are very sensitive to steroids.
- ▶ Cats or horses suspected of having viral (eg, herpes) or fungal eye infections.
- ▶ Animals with eye ulcers.

This drug should be used **WITH CAUTION** in patients that:

- ▶ Are diabetic.
- ▶ Have glaucoma.
- ▶ Are pregnant or nursing.

If your pet has any of these conditions or signs, talk to your veterinarian about the potential risks versus benefits of using this medication.

What are the side effects of this medication?

Side effects that usually are not serious include:

- ▶ Mild burning, stinging, irritation, or redness of the eyes.
- ▶ Drooping of the eyelid.

If any of these are severe, worsen, or continue to be a problem, contact your veterinarian as soon as possible to avoid complications.

Side effects that may be serious or indicate a serious problem:

- ▶ Difficulty breathing or swelling of the throat, which can indicate a severe allergic reaction to this drug.
- ▶ Signs of an eye infection such as squinting, yellow or green discharge, and sensitivity to light.
- ▶ Failure of eye injuries to heal.
- ▶ Loss of blood sugar control in diabetic animals.

If your pet has any of these conditions or signs, contact your veterinarian immediately.

If my pet gets too much of this medication (an overdose), what should I do?

When used as an eye medication, overdoses are not likely, but side effects or toxic effects could occur if your animal eats the medication. If this happens, contact your veterinarian or an animal poison

control center for further advice. Animal poison control centers that are open 24 hours a day include **ASPCA Animal Poison Control Center** (888-426-4435) and **Pet Poison HELPLINE** (855-764-7661); a consultation fee may be charged for these services.

How should this medication be given?

Give this medication exactly as your veterinarian has prescribed. It's a good idea to always check the prescription label to be sure you are giving the drug correctly.

- ▶ Some forms of this drug must be shaken well prior to using. Read the instructions provided by your veterinarian or pharmacist.
- ▶ Wash your hands before administering this medication.
- ▶ Do not touch the dropper tip to your animal's eye or any other surface to prevent contamination.
- ▶ If any residue is left on your animal's face after giving the eye drops, gently wipe it off with a damp cloth or tissue.
- ▶ If you are administering more than one eye medication to your animal, wait 5 minutes between each medication before giving the next one. Use eye drops before eye ointments to allow the drops to absorb into the eye.
- ▶ If you are using this medication as an eye solution for your horse, your veterinarian may have implanted a special eye catheter (subpalpebral lavage [SPL] catheter). Use this catheter exactly as your veterinarian has prescribed. Only use air to flush the medications into the eye once injected into the catheter.
- ▶ If you are having difficulty applying the medication or your animal does not accept the treatment, contact your veterinarian or pharmacist for tips to help with administration and reducing the stress of medication time.
- ▶ This medication can be given for various lengths of time. Be sure you understand how long your veterinarian wants you to continue giving this medication. Prescription refills may be necessary before the therapy will be complete. Before stopping this medication, talk to your veterinarian, as there may be important reasons to continue its use.

What should I do if I miss giving a dose of this medication?

If you miss a dose, give it when you remember. However, if it is close to the time for the next dose, skip the dose you missed and give it

at the next scheduled time. After that, return to the regular dosing schedule. Do not double-up or give extra doses.

How should I store this medication?

- ▶ Store this medication in the original prescription bottle or an approved dosage reminder (ie, pill minder) container at room temperature and protected from light.
- ▶ If your veterinarian or pharmacist has made (compounded) a special formulation for your animal, follow the storage recommendations and expiration date for the product.
- ▶ Keep away from children and other animals.

Can handling this medication be hazardous to me, my family, or other pets?

There are no specific precautions required when handling this medication unless you are allergic to it or other drugs like it (eg, dexamethasone), but it is always a good idea to wash your hands after handling any medication.

How should I dispose of this medication if I don't use it all?

- ▶ Do not flush this medication down the toilet or wash it down the sink. If a community drug "take-back" program is available, use this option. If there is no take-back program, mix the drug with coffee grounds or cat litter (to make it undesirable to children and animals and unrecognizable to people who might go through your trash), place the mixture in a sealable plastic bag to keep it from leaking out, and throw the bag out with the regular trash.
- ▶ Do not save left over medication for future use or give it to others to use.

What other information is important for this medication?

- ▶ Use of this drug may not be allowed in certain animal competitions. Check rules and regulations before entering your animal in a competition while this medication is being administered.

If you have any other questions or concerns about this medication, contact your veterinarian or pharmacist.