

**DERMALONE - nystatin, neomycin sulfate, thioestrepton and triamcinolone acetonide ointment**  
**Vedco**

-----  
**Nystatin, Neomycin Sulfate, Thioestrepton and Triamcinolone Acetonide Ointment**

**For Use Only in Dogs and Cats**  
**FDA.**

**NADA 140-810, Approved by**

**CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.**

**DESCRIPTION:**

Nystatin, neomycin sulfate, thioestrepton and triamcinolone acetonide ointment is an ointment in a non-irritating vehicle, a polyethylene and mineral oil gel base.

Each mL contains:

nystatin 100,000 units

neomycin sulfate (equivalent to neomycin base) 2.5 mg

thioestrepton 2,500 units

triamcinolone acetonide 1.0 mg

Nystatin, neomycin sulfate, thioestrepton and triamcinolone acetonide ointment is supplied in tubes of 1/4 fl.oz. (7.5 mL.), 1/2 fl. oz. (15 mL.), and 1 fl. oz. (30 mL.), each with an elongated tip for easy application and in dispensing bottles of 8 fl. oz. (240 mL.). The preparation is intended for local therapy in a variety of cutaneous disorders of cats and dogs; it is especially useful in disorders caused, complicated, or threatened by bacterial and/or candidal (monilia) infections.

**ACTIONS:**

By virtue of its four active ingredients, the ointment provides four basic therapeutic effects: anti-inflammatory, antipruritic, antifungal and antibacterial. Triamcinolone acetonide is a potent synthetic corticosteroid providing rapid and prolonged symptomatic relief on topical administration.

Inflammation, edema, and pruritis promptly subside and lesions are permitted to heal. Nystatin is the first well tolerated anti-fungal antibiotic of dependable efficacy for the treatment of cutaneous infections caused by Candida albicans (monilia). Nystatin is fungistatic in vitro against a variety of yeast and yeast-like fungi including many fungi pathogenic to animals. No appreciable activity is exhibited against bacteria. Thioestrepton has a high order of activity against gram-positive organisms, including many which are resistant to other antibiotics; neomycin exerts antimicrobial action against a wide range of gram-positive and gram-negative bacteria. Together they provide comprehensive therapy against those organisms responsible for most superficial bacterial infections.

**INDICATIONS:**

Nystatin, neomycin sulfate, thioestrepton and triamcinolone acetonide ointment is particularly useful in the treatment of acute and chronic otitis of varied etiologies, in interdigital cysts in cats and dogs, and in anal gland infections in dogs. The preparation is also indicated in the management of dermatologic disorders characterized by inflammation and dry or exudative dermatitis, particularly those caused, complicated, or threatened by bacterial or candidal (Candida albicans) infections. It is also of value in eczematous dermatitis; contact dermatitis, and seborrheic dermatitis; and as an adjunct in the treatment of dermatitis due to parasitic infestation.

**WARNINGS:**

Clinical and experimental data have demonstrated that corticosteroids administered orally or by injection to animals may induce the first stage of parturition if used during the last trimester of pregnancy and may precipitate premature parturition followed by dystocia, fetal death, retained placenta, and metritis.

Additionally, corticosteroids administered to dogs, rabbits and rodents during pregnancy have resulted in cleft palate in the offspring. In dogs, other congenital anomalies have resulted; deformed forelegs,

phocomelia, and anasarca.

### **PRECAUTIONS:**

Nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment is not intended for the treatment of deep abscesses or deep-seated infections such as inflammation of the lymphatic vessels. Parenteral antibiotic therapy is indicated in these infections.

Nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment has been extremely well tolerated. Cutaneous reactions attributable to its use have been extremely rare. The occurrence of systemic reactions is rarely a problem with topical administration.

There is some evidence that corticosteroids can be absorbed after topical application and cause systemic effects. Therefore, an animal receiving nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment therapy should be observed closely for signs such as polydipsia, polyuria, and increased weight gain. Nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment is not generally recommended for the treatment of deep or puncture wounds or serious burns.

Sensitivity to neomycin may occur. If redness, irritation or swelling persists or increases, discontinue use. Do not use if pus is present since the drug may allow the infection to spread. Keep this and all medications out of the reach of children.

Avoid ingestion. Oral or parenteral use of corticosteroids (depending on dose, duration of use, and specific steroid) may result in inhibition of endogenous steroid production following drug withdrawal.

### **SIDE EFFECTS:**

SAP and SGPT (ALT) enzyme elevations, polydipsia/polyuria, vomiting, and diarrhea (occasionally bloody) have been observed following parenteral or systemic use of synthetic corticosteroids in dogs.

Cushing's syndrome has been reported in association with prolonged or repeated steroid therapy in dogs.

Temporary hearing loss has been reported in conjunction with treatment of otitis with products containing corticosteroids. However, regression usually occurred following withdrawal of the drug. If hearing dysfunction is noted during the course of treatment with nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment, discontinue its use.

### **CAUTION:**

Before instilling any medication into the ear, examine the external ear canal thoroughly to be certain the tympanic membrane is not ruptured in order to avoid the possibility of transmitting infection to the middle ear as well as damaging the cochlea or vestibular apparatus from prolonged contact. If hearing or vestibular dysfunction is noted during the course of treatment, discontinue the use of nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment.

### **DOSAGE AND ADMINISTRATION:**

Frequency of administration is dependent on the severity of the condition. For mild inflammations, application may range from once daily to once a week; for severe conditions nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment may be applied as often as two to three times daily, if necessary. Frequency of treatment may be decreased as improvement occurs.

Wear gloves during the administration of the ointment or wash hands immediately after application.

**OTITIS:** Clean ear canal of impacted cerumen. Inspect canal and remove any foreign bodies such as grass awns, ticks, etc. Instill three to five drops of nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment.

Preliminary use of a local anesthetic such as Proparacaine Hydrochloride Ophthalmic Solution may be advisable.

**INFECTED ANAL GLANDS, CYSTIC AREAS, ETC:** Drain gland or cyst and then fill with nystatin, neomycin sulfate, thioestrepton and triamcinolone acetonide ointment.

**OTHER DERMATOLOGIC DISORDERS:** Clean affected areas, removing any encrusted discharge or exudate. Apply nystatin, neomycin sulfate, thioestrepton and triamcinolone acetonide ointment sparingly in a thin film.

**STORAGE:**

240 mL bottle: Do not store above 86°F. 7.5 mL, 15 mL, 30 mL tubers: Store at room temperature; avoid excessive heat (104°F).

**Manufactured by Med-Pharmex, Inc.  
Pomona, CA 91767**

Revised 10/98

The image shows the front and back panels of a Dermalone Ointment tube. The front panel (left) is white with black text. At the top, it lists the NDC number (50989-383-66) and the product name 'DERMALONE™ OINTMENT'. Below that, it lists the active ingredients: 'NYSTATIN-NEOMYCIN SULFATE-THIOSTREPTON-TRIAMCINOLONE ACETONIDE OINTMENT'. The manufacturer information is 'Manufactured For VEDCO, INC., St. Joseph, MO 64507'. At the bottom, it lists the contents of each mL: Nystatin (100,000 Units), Neomycin Sulfate (Equivalent to Neomycin Base) (2.5 mg), Thioestrepton (2,500 Units), and Triamcinolone Acetonide (1 mg). It also notes the base is a polyethylene and mineral oil gel.

The back panel (right) is light gray with a blue diagonal band. At the top, it has a 'CAUTION' statement: 'Federal law restricts this drug to use by or on the order of a licensed veterinarian.' The blue band features the 'VEDCO' logo. Below the band, it states 'Net Contents: 7.5 mL (1/4 Fl. oz.)' and 'NADA 140-810 Approved by FDA'. At the bottom, it provides usage instructions: 'For use in dogs and cats only', 'Dosage and administration: Refer to insert. Store at room temperature. Avoid excessive heat. (104°)', and 'Exp. date and lot no. on crimp'.

NDC 50989-383-08

# DERMALONE™ OINTMENT

NYSTATIN-NEOMYCIN  
SULFATE-THIOSTREPTON-  
TRIAMCINOLONE ACETONIDE  
OINTMENT

Manufactured For  
**VEDCO, INC.**  
St. Joseph, MO 64507

## DERMALONE™ OINTMENT

NYSTATIN-NEOMYCIN SULFATE-  
THIOSTREPTON-TRIAMCINOLONE  
ACETONIDE OINTMENT.

Each mL Contains:

Nystatin 100,000 Units  
Neomycin Sulfate  
(Equivalent to Neomycin  
Base) 2.5 mg.  
Thiostrepton 2,500 Units  
Triamcinolone  
Acetonide 1 mg.  
In a polyethylene and  
mineral oil gel base

**CAUTION:** Federal law restricts  
this drug to use by or on the order  
of a licensed veterinarian.



Net Contents: 15 mL (1/2 Fl. oz.)  
NADA 140-810 Approved by FDA

**For use in dogs  
and cats only**

**Dosage and administration:**  
Refer to insert. Store at  
room temperature. Avoid  
excessive heat. (104°F)

**Exp. date and lot no. on crimp**

NDC 50989-383-10

# DERMALONE™ OINTMENT

NYSTATIN-NEOMYCIN  
SULFATE-THIOSTREPTON-  
TRIAMCINOLONE ACETONIDE  
OINTMENT

Manufactured For  
**VEDCO, INC.**  
St. Joseph, MO 64507

## DERMALONE™ OINTMENT

NYSTATIN-NEOMYCIN SULFATE-  
THIOSTREPTON-TRIAMCINOLONE  
ACETONIDE OINTMENT

Each mL Contains:

Nystatin . . . 100,000 Units  
Neomycin Sulfate  
(Equivalent to Neomycin  
Base) . . . . . 2.5 mg  
Thiostrepton 2,500 Units  
Triamcinolone  
Acetonide . . . . . 1 mg  
In a polyethylene and  
mineral oil gel base.

**CAUTION:** Federal law restricts  
this drug to use by or on the order  
of a licensed veterinarian.



# VEDCO

Net Contents: 30 mL (1 Fl. oz.)  
NADA 140-810 Approved by FDA

**For use in dogs and  
cats only**

**Dosage and administration:  
Refer to insert. Store at  
room temperature. Avoid  
excessive heat (104°F.)**

**Expiration date and lot number on crimp.**



NDC 50989-383-24

# DERMALONE OINTMENT™

NYSTATIN-NEOMYCIN SULFATE-  
THIOSTREPTON-TRIAMCINOLONE  
ACETONIDE OINTMENT

Net Contents: 240 mL (8 fl.oz.)

# VEDCO

**CAUTION:**  
Federal law  
restricts this drug  
to use by or on the order  
of a licensed veterinarian.

Manufactured for  
**VEDCO, INC.**  
St. Joseph, MO 64507

NADA 140-810 Approved by FDA

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

Each mL Contains:

Nystatin 100,000 Units

Neomycin Sulfate (equivalent  
to neomycin base) 2.5 mg.

Thiostrepton 2,500 Units

Triamcinolone Acetonide 1 mg.  
In a polyethylene and mineral  
oil gel base.

Shake well before using.

**For use in dogs and cats  
only.**

Dosage and administration:  
Refer to insert.

Do not store above 86° F.

Lot #/Exp. Date

## DERMALONE

nystatin, neomycin sulfate, thiostrepton and triamcinolone acetonide ointment

### Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:50989-383
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Nystatin (UNII: BDF1O1C72E) (Nystatin - UNII:BDF1O1C72E)	Nystatin	100000 [iU] in 1 mL
Neomycin Sulfate (UNII: 057Y626693) (Neomycin - UNII:I16QD7X297)	Neomycin	2.5 mg in 1 mL
Thiostrepton (UNII: HR4S203Y18) (Thiostrepton - UNII:HR4S203Y18)	Thiostrepton	2500 [iU] in 1 mL
Triamcinolone acetonide (UNII: F446C597KA) (Triamcinolone acetonide - UNII:F446C597KA)	Triamcinolone acetonide	1 mg in 1 mL

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50989-383-66	7.5 mL in 1 TUBE		
2	NDC:50989-383-08	15 mL in 1 TUBE		
3	NDC:50989-383-10	30 mL in 1 TUBE		
4	NDC:50989-383-24	240 mL in 1 BOTTLE, PLASTIC		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA140810	10/01/1998	

**Labeler** - Vedco (021634266)**Registrant** - Med-Pharmex, Inc. (025353699)**Establishment**

Name	Address	ID/FEI	Business Operations
Med-Pharmex, Inc.		025353699	manufacture

Revised: 10/1998

Vedco