

CHLORHEXIDINE +PS SHAMPOO with CLIMBAZOLE



Antiseptic lathering solution with Chlorhexidine, Climbazole and Phytosphingosine

TECHNICAL SPECIFICATIONS:

Lathering solution formulated with gentle hypoallergenic cleansing bases
Phytosphingosine – salicyloyl (0.05%) – Climbazole 0.5% - Lipacide® C8G (2.5%) - Chlorhexidine (3%)

INGREDIENTS:

Aqua, Sulfobetaine, Chlorhexidine Digluconate, Climbazole, Decylpolyglucoside, Capryloyl Glycine, Peg-120, Methyl Glucose Dioleate, Sodium Hydroxide, Triethanolamine, Fragrance, Salicyloyl Phytosphingosine, Citric Acide.

PROPERTIES:

Chlorhexidine has a strong bactericidal and bacteriostatic activity on the Gram positive and Gram negative bacteria. Climbazole is an imidazole compound active on most fungi and yeasts (*Malassezia pachydermatis*).

Phytosphingosine is a pro-ceramide that helps restore the skin barrier, and also has anti-inflammatory properties.

Both the antibacterial and antifungal activities of the lipo-amino-acid complement those of chlorhexidine. The Lipacid also acts as a moisturizing and restructuring agent by allowing a lipo-protidic film to deposit on the skin. This long-lasting film resists rinsing, and thus protects the skin from irritations and loss of moisture.

The formula of DOUXO® Chlorhexidine is suitable for frequent use.

DIRECTIONS FOR USE:

External use only. Wet the animal with warm water and apply DOUXO® Calm Shampoo extensively on the fur to create a rich lather. Massage the shampoo into the wet hair coat, lather freely. Allow to remain on the hair coat for 5 to 10 minutes before rinsing thoroughly with clean water. Repeat if needed. The veterinarian's recommendations should be observed.

It is generally recommended to shampoo 2 to 3 times weekly, for 1 to 2 weeks. Then, for effective control of the condition, can be used in shift with DOUXO® Chlorhexidine Micro-emulsion Spray and/or Douxo Chlorhexidine +Climbazole Mousse.

SPECIES:

Dogs and cats: topical adjuvant management of superficial and severe skin conditions that are responsive to chlorhexidine and climbazole.

PRECAUTIONS:

Do not swallow. Avoid contact with the eyes. If undue skin irritation develops or increases, discontinue use and contact a veterinarian. Keep out of reach of children. To facilitate the use of the product, bottles can be stored upside down.

STORAGE:

Store at controlled room temperature.

PRESENTATION:

Bottles of 200 ml (6.8 fl. oz.), 500 ml (16.9 fl. oz.) and 3L (101.4 fl. oz)

